



(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 September 2003 (25.09.2003)

PCT

(10) International Publication Number  
WO 03/078627 A2

- (51) International Patent Classification: C12N 15/10
- (21) International Application Number: PCT/DK03/00177
- (22) International Filing Date: 14 March 2003 (14.03.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
PA 2002 0415 15 March 2002 (15.03.2002) DK  
60364 056 15 March 2002 (15.03.2002) US  
PCT/DK 02/00419 20 June 2002 (20.06.2002) DK  
10175 539 20 June 2002 (20.06.2002) US  
60434 439 19 December 2002 (19.12.2002) US
- (71) Applicant (for all designated States except US): NUEVO-  
LUTION A/S [DK/DK]; Rønnegade 8, 5th floor, DK-2100  
Copenhagen Ø (DK).
- (72) Inventors: and
- (73) Inventors/Applicants (for US only): GOULIAEV, Alex,  
Haahr [DK/DK]; Brøndsted 223, DK-3670 Veksø Sjeel-  
land (DK); PEDERSEN, Henrik [DK/DK]; Frodesvej  
24, DK-2880 Bagsvaerd (DK); THISTED, Thomas  
[DK/DK]; Fjødskretzen 14, DK-3600 Frederikssund  
(DK); LUNDØRE, Mikkel, Dybro [DK/DK]; Charlotte  
Munksvøj 31, 2. tv., DK-2400 København NV (DK);  
SAMSS, Christian [DK/DK]; Jakob Dammeferds Vej 4 A,
- Published:  
— without international search report and to be republished  
upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(34) Title: A BUILDING BLOCK CAPABLE OF FUNCTIONAL ENTITY TRANSFER TO NUCLEOPHIL

(57) Abstract: A building block having the dual capabilities of transferring the genetic information e.g. by recognising an encoding element and transferring a functional entity to a recipient reactive group is disclosed. The building block can be designed with an adjustable transferability taking into account the components of the building block. The building block may be used in the generation of a single complex or libraries of different complexes, wherein the complex comprises an encoded molecule linked to an encoding element. Libraries of complexes are useful in the quest for pharmaceutically active compounds.

Title

A BUILDING BLOCK CAPABLE OF FUNCTIONAL ENTITY TRANSFER TO NU-  
CLEOPHIL

5 Technical Field of the Invention

The present invention relates to a building block comprising a complementing ele-  
ment and precursor for a functional entity. The building block is designed to transfer  
the functional entity with an adjustable efficiency to a recipient reactive group upon  
recognition between the complementing element and an encoding element associ-  
ated with the reactive group. The invention also relates to a linkage between the  
functional entity and the complementing element as well as a method for transferring  
a functional entity to recipient reactive group.

Background

15 The transfer of a chemical entity from one mono-, di- or oligonucleotide to another  
has been considered in the prior art. Thus, N. M. Chung *et al.* (Biochim. Biophys.  
Acta, 1971, 228:536-543) used a poly(U) template to catalyse the transfer of an ace-  
tyl group from 3'-O-acetyladenosine to the 5'-OH of adenosine. The reverse transfer,  
i.e. the transfer of the acetyl group from a 5'-O-acetyladenosine to a 3'-OH group of  
another adenosine, was also demonstrated.

25 Walder *et al.* Proc. Natl. Acad. Sci. USA, 1979, 76, 51-55 suggest a synthetic pro-  
cedure for peptide synthesis. The synthesis involves the transfer of nascent immobi-  
lized polypeptide attached to an oligonucleotide strand to a precursor amino acid  
attached to an oligonucleotide. The transfer comprises the chemical attack of the  
amino group of the amino acid precursor on the substitution labile peptidyl ester,  
which in turn results in an acyl transfer. It is suggested to attach the amino acid pre-  
cursor to the 5' end of an oligonucleotide with a thiol ester linkage.

30 The transfer of a peptide from one oligonucleotide to another using a template is  
disclosed in Bruick RK *et al.* Chemistry & Biology, 1996, 3:49-56. The carboxy ter-  
minal of the peptide is initially converted to a thioester group and subsequently  
transformed to an activated thioester upon incubation with Ellman's reagent. The  
activated thioester is reacted with a first oligo, which is 5'-thiol-terminated, resulting  
in the formation of a thio-ester linked intermediate. The first oligonucleotide and a

second oligonucleotide having a 3' amino group is aligned on a template such that the thioester group and the amino group are positioned in close proximity and a reaction is effected resulting in a coupling of the peptide to the second oligonucleotide through an amide bond.


5

The prior art building blocks and methods for transfer have a relatively poor transfer efficiency. Therefore, in an aspect of the present invention an oligonucleotide conjugated to a transferable chemical moiety via a linker is provided, which has an increased ability to transfer a functional entity.

10

### Summary of the Invention

The present invention relates to a building block of the general formula




15

capable of transferring a functional entity (FE) to a recipient reactive group, wherein the lower horizontal line is a **Complementing Element** identifying the functional entity and the vertical line between the complementing element and the S atom is a **Spacer**.

20

Preferably the spacer is a valence bond,  $C_1-C_6$  alkylene-A-,  $C_1-C_6$  alkenylene-A-,  $C_2-C_6$  alkynylene-A-, or



said spacer optionally being connected through A to a moiety selected from




25

$-(CH_2)_n-S-S-(CH_2)_m-B-$  where A is a valence bond,  $-C(O)NR^1-$ ,  $-NR^1-$ ,  $-O-$ ,  $-S-$ , or  $-C(O)-O-$ ; B is a valence bond,  $-O-$ ,  $-S-$ ,  $-NR^1-$  or  $-C(O)NR^1-$  and connects to the S atom of the carrier;  $R^1$  is selected independently from H,  $C_1-C_6$  alkyl,  $C_2-C_7$  cycloalkyl,  $C_1-C_6$  alkenylene-aryl, or aryl substituted with 0-5 halogen atoms selected from -F, -Cl, -Br and -I; and n and m independently are integers ranging from 1 to 10.



5

In one aspect of the invention the **Spacer** is  $C_1-C_6$  alkylene-A-,  $C_1-C_6$  alkenylene-A-,  $C_2-C_6$  alkynylene-A-, or



10


said spacer optionally being connected through A to a moiety selected from



where A is  $-C(O)NR^1-$ , or  $-S-$ ; B is  $-S-$ ,  $-NR^1-$  or  $-C(O)NR^1-$  and connects to S-C-connecting group;  $R^1$  is selected independently from H,  $C_1-C_6$  alkyl,  $C_1-C_6$  alkenylene-aryl, or aryl; and n and m independently are integers ranging from 1 to 6.


15

Preferably the **Spacer** is -A-, a group  $C_1-C_6$  alkylene-A-,  $C_2-C_6$  alkenylene-A-, or  $C_2-C_6$  alkynylene-A- optionally substituted with 1 to 3 hydroxy groups, or



20

said spacer being connected through A to a linker selected from



where A is a valence bond,  $-NR^2-$ ,  $-C(O)NR^2-$ ,  $-NR^2-C(O)-$ ,  $-O-$ ,  $-S-$ ,  $-C(O)-O-$  or  $OP(=O)(O^-)-O-$ ; B is a valence bond,  $-O-$ ,  $-S-$ ,  $-NR^2-$ ,  $-C(O)-$  or  $-C(O)NR^2$ , and connects to S-C-connecting group;  $R^2$  is selected independently from H,  $C_1-C_6$  alkyl,

25



R<sup>8</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, aryl or C<sub>1</sub>-C<sub>6</sub> alkylene-aryl substituted with 0-3 substituents independently selected from -F, -Cl, -NO<sub>2</sub>, -R<sup>3</sup>, -OR<sup>3</sup>, -SR<sup>3</sup>, R<sup>9</sup> is O, -F, -Cl, -Br, -I, -CN, -NO<sub>2</sub>, -OR<sup>9</sup>, -NR<sup>9</sup>, -NR<sup>9</sup>-C(O)R<sup>9</sup>, -NR<sup>9</sup>-C(O)OR<sup>9</sup>, -SR<sup>9</sup>, -S(O)R<sup>9</sup>, -S(O)<sub>2</sub>R<sup>9</sup>, -COOR<sup>9</sup>, -C(O)NR<sup>9</sup> and -S(O)<sub>2</sub>NR<sup>9</sup>.

5

In a certain aspect of the invention, R is H or selected among the group consisting of a C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkenyl, C<sub>3</sub>-C<sub>7</sub> cycloalkynyl, aryl, and heteroaryl, said group being substituted with 0-3 R<sup>5</sup> and 0-3 R<sup>6</sup>, or selected among the group consisting of C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>-C(O)R<sup>8</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>-C(O)OR<sup>8</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>-C(O)R<sup>8</sup>, and C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>-C(O)OR<sup>8</sup> substituted with 0-3 R<sup>6</sup>.

10

Suitably, R is H or selected among the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkenyl, C<sub>3</sub>-C<sub>7</sub> cycloalkynyl, aryl, and heteroaryl, said group being substituted with 0-3 R<sup>5</sup> and 0-3 R<sup>6</sup>.

15

In some aspects of the invention it is preferred that R is selected among the group consisting of C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>-C(O)R<sup>8</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>-C(O)OR<sup>8</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>-C(O)R<sup>8</sup>, and C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>-C(O)OR<sup>8</sup> substituted with 0-3 R<sup>6</sup>.

20

In the present description and claims, the direction of connections between the various components of a building block should be read left to right. For example a spacer is connected to a complementing element through the atom on the left and to the sulphur atom (or alternatively the group A) through the atom on the right hand side.

25

The term "C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl" as used herein refers to a radical of totally saturated heterocycle like a cyclic hydrocarbon containing one or more heteroatoms selected from nitrogen, oxygen, phosphor, boron and sulphur independently in the cycle such as pyrrolidine (1- pyrrolidine; 2- pyrrolidine; 3- pyrrolidine; 4- pyrrolidine; 5- pyrrolidine); pyrazolidine (1- pyrazolidine; 2- pyrazolidine; 3- pyrazolidine; 4- pyrazolidine; 5- pyrazolidine); imidazolidine (1- imidazolidine; 2- imidazolidine; 3- imidazolidine; 4- imidazolidine; 5- imidazolidine); thiazolidine (2- thiazolidine; 3- thiazolidine; 4- thiazolidine; 5- thiazolidine).

35

zolidine; 3- thiazolidine; 4- thiazolidine; 5- thiazolidine); piperidine (1- piperidine; 2- piperidine; 3- piperidine; 4- piperidine; 5- piperidine; 6- piperidine); piperazine (1- piperazine; 2- piperazine; 3- piperazine; 4- piperazine; 5- piperazine; 6- piperazine); morpholine (2- morpholine; 3- morpholine; 4- morpholine; 5- morpholine; 6- morpholine); thiomorpholine (2- thiomorpholine; 3- thiomorpholine; 4- thiomorpholine; 5- thiomorpholine; 6- thiomorpholine); 1,2-oxathiolane (3-(1,2-oxathiolane); 4-(1,2-oxathiolane); 5-(1,2-oxathiolane); 1,3-dioxolane (2-(1,3-dioxolane); 4-(1,3-dioxolane); 5-(1,3-dioxolane); tetrahydropyran; 6- tetrahydropyran; 3-tetrahydropyran; 4-tetrahydropyran; 5-tetrahydropyran; 6- tetrahydropyran); hexahydropyridazine (1-(hexahydropyridazine); 2- (hexahydropyridazine); 3-(hexahydropyridazine); 4-(hexahydropyridazine); 5- (hexahydropyridazine); 6-(hexahydropyridazine)), [1,3,2]dioxaborolane, [1,3,6,2]dioxazaborocane

5

10

The term "aryl" as used herein includes carbocyclic aromatic ring systems of 5-7 carbon atoms. Aryl is also intended to include the partially hydrogenated derivatives of the carbocyclic systems as well as up to four fused aromatic- or partially hydrogenated rings, each ring comprising 5-7 carbon atoms.

15

The term "heteroaryl" as used herein includes heterocyclic unsaturated ring systems containing, in addition to 2-18 carbon atoms, one or more heteroatoms selected from nitrogen, oxygen and sulphur such as furyl, thienyl, pyrrolyl, heteroaryl is also intended to include the partially hydrogenated derivatives of the heterocyclic systems enumerated below.

20

The terms "aryl" and "heteroaryl" as used herein refers to an aryl which can be optionally substituted or a heteroaryl which can be optionally substituted and includes phenyl, biphenyl, indenyl, naphthyl (1-naphthyl, 2-naphthyl), N-hydroxytetrazolyl, N-hydroxytriazolyl, N-hydroxyimidazolyl, anthracenyl (1-anthracenyl, 2-anthracenyl, 3-anthracenyl), thiophenyl (2-thienyl, 3-thienyl), furyl (2-furyl, 3-furyl), indolyl, oxadiazolyl, isoxazolyl, quinazolyl, fluorenyl, xanthenyl, isoindanyl, benzhydryl, acridinyl, thiazolyl, pyrrolyl (2-pyrrolyl), pyrazolyl (3-pyrazolyl), imidazolyl (1-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl), triazolyl (1,2,3-triazol-1-yl, 1,2,3-triazol-2-yl, 1,2,3-triazol-4-yl, 1,2,4-triazol-3-yl), oxazolyl (2-oxazolyl, 4-oxazolyl, 5-oxazolyl), thiazolyl (2-thiazolyl, 4-thiazolyl, 5-thiazolyl), pyridyl (2-pyridyl, 3-pyridyl, 4-pyridyl), pyrimidinyl (2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, 6-pyrimidinyl), pyrazinyl, pyridazinyl (3-pyridazinyl, 4-pyridazinyl, 5-pyridazinyl), quinolyl (2-quinolyl, 3-quinolyl, 4-quinolyl, 5-quinolyl, 6-

25

30

35


quinoilyl, 7-quinoly, 8-quinoly), isquinoly (1-isquinoly, 3-isquinoly, 4-isquinoly, 5-isquinoly, 6-isquinoly, 7-isquinoly, 8-isquinoly), benzo[b]furanyl (2-benzo[b]furanyl, 3-benzo[b]furanyl, 4-benzo[b]furanyl, 5-benzo[b]furanyl, 6-benzo[b]furanyl, 7-benzo[b]furanyl), 2,3-dihydro-benzo[b]furanyl (2-(2,3-dihydro-benzo[b]furanyl), 3-(2,3-dihydro-benzo[b]furanyl), 4-(2,3-dihydro-benzo[b]furanyl), 5-(2,3-dihydro-benzo[b]furanyl), 6-(2,3-dihydro-benzo[b]furanyl), 7-(2,3-dihydro-benzo[b]furanyl), benzo[b]thiophenyl (2-benzo[b]thiophenyl, 3-benzo[b]thiophenyl, 4-benzo[b]thiophenyl, 5-benzo[b]thiophenyl, 6-benzo[b]thiophenyl, 7-benzo[b]thiophenyl), 2,3-dihydro-benzo[b]thiophenyl (2-(2,3-dihydro-benzo[b]thiophenyl), 3-(2,3-dihydro-benzo[b]thiophenyl), 4-(2,3-dihydro-benzo[b]thiophenyl), 5-(2,3-dihydro-benzo[b]thiophenyl), 6-(2,3-dihydro-benzo[b]thiophenyl), 7-(2,3-dihydro-benzo[b]thiophenyl), indolyl (1-indolyl, 2-indolyl, 3-indolyl, 4-indolyl, 5-indolyl, 6-indolyl, 7-indolyl), indazole (1-indazolyl, 3-indazolyl, 4-indazolyl, 5-indazolyl, 6-indazolyl, 7-indazolyl), benzimidazolyl (1-benzimidazolyl, 2-benzimidazolyl, 4-benzimidazolyl, 5-benzimidazolyl, 6-benzimidazolyl, 7-benzimidazolyl, 8-benzimidazolyl), benzoxazolyl (1-benzoxazolyl, 2-benzoxazolyl), benzothiazolyl (1-benzothiazolyl, 2-benzothiazolyl, 4-benzothiazolyl, 5-benzothiazolyl, 6-benzothiazolyl, 7-benzothiazolyl), carbazolyl (1-carbazolyl, 2-carbazolyl, 3-carbazolyl, 4-carbazolyl), 5H-dibenz[b]flazepine (5H-dibenz[b]flazepin-1-yl, 5H-dibenz[b]flazepine-2-yl, 5H-dibenz[b]flazepine-3-yl, 5H-dibenz[b]flazepine-4-yl, 5H-dibenz[b]flazepine-5-yl), 10,11-dihydro-5H-dibenz[b]flazepine (10,11-dihydro-5H-dibenz[b]flazepine-1-yl, 10,11-dihydro-5H-dibenz[b]flazepine-2-yl, 10,11-dihydro-5H-dibenz[b]flazepine-3-yl, 10,11-dihydro-5H-dibenz[b]flazepine-4-yl, 10,11-dihydro-5H-dibenz[b]flazepine-5-yl).

The Functional Entity carries elements used to interact with host molecules and optionally reactive elements allowing further elaboration of an encoded molecule of a library. Interaction with host molecules like enzymes, receptors and polymers is typically mediated through van der Waals interactions, polar- and ionic interactions and pi-stacking effects. Substituents mediating said effects may be masked by methods known to an individual skilled in the art (Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*; 3rd ed.; John Wiley & Sons: New York, 1999.) to avoid undesired interactions or reactions during the preparation of the individual building blocks and during library synthesis. Analogously, reactive elements may be masked by suitably selected protection groups. It is appreciated by one skilled in the

art that by suitable protection, a functional entity may carry a wide range of substituents.


The Functional Entity may be a masked Functional Entity that is incorporated into an encoded molecule. After incorporation, reactive elements of the Functional Entity may be revealed by un-masking allowing further synthetic operations. Finally, elements mediating recognition of host molecules may be un-masked.

The function of the carrier




is to provide for the transferability of the functional entity, playing the role of a leaving group.

The spacer serves to distance the functional entity to be transferred from the bulky complementing element. Thus, the identity of the spacer is not crucial for the function of the building block. It may be desired to have a spacer which can be cleaved by light. In this occasion, the spacer is provided with e.g. the group



In the event an increased hydrophilicity is desired the spacer may be provided with a polyethylene glycol part of the general formula:



The spacer in conjunction with the carrier makes up a cleavable linker, which links the complementing element to the functional entity.

In a preferred embodiment, the complementing element serves the function of transferring genetic information e.g. by recognising a coding element. The recognition implies that the two parts are capable of interacting in order to assemble a complementing element - coding element complex. In the biotechnological field a variety of interacting molecular parts are known which can be used according to the invention. Examples include, but are not restricted to protein-protein interactions, protein-polysaccharide interactions, RNA-protein interactions, DNA-DNA interactions, DNA-RNA interactions, RNA-RNA interactions, biotin-streptavidin interactions, enzyme-ligand interactions, antibody-ligand interaction, protein-ligand interaction, ect.

10

The interaction between the complementing element and coding element may result in a strong or a weak bonding. If a covalent bond is formed between the parties of the affinity pair the binding between the parts can be regarded as strong, whereas the establishment of hydrogen bondings, interactions between hydrophobic domains, and metal chelation in general results in weaker bonding. In general relatively weak bonding is preferred. In a preferred aspect of the invention, the complementing element is capable of reversible interacting with the coding element so as to provide for an attachment or detachment of the parts in accordance with the changing conditions of the media.

15

20

In a preferred aspect of the invention, the interaction is based on nucleotides, i.e. the complementing element is a nucleic acid. Preferably, the complementing element is a sequence of nucleotides and the coding element is a sequence of nucleotides capable of hybridising to the complementing element. The sequence of nucleotides carries a series of nucleobases on a backbone. The nucleobases may be any chemical entity able to be specifically recognized by a complementing entity. The nucleobases are usually selected from the natural nucleobases (adenine, guanine, uracil, thymine, and cytosine) but also the other nucleobases obeying the Watson-Crick hydrogen-bonding rules may be used, such as the synthetic nucleobases disclosed in US 6,037,120. Examples of natural and non-natural nucleobases able to perform a specific pairing are shown in Figure 2. The backbone of the sequence of nucleotides may be any backbone able to aggregate the nucleobases in a sequence. Examples of backbones are shown in figure 4. In some aspects of the invention the addition of non-specific nucleobases to the complementing element is advantageous, figure 3.

35

SUBSTITUTE SHEET (RULE 26)

The coding element can be an oligonucleotide having nucleobases which complement and is specifically recognised by the complementing element, i.e. in the event the complementing element contains cytosine, the coding element part contains guanine and visa versa, and in the event the complementing element contains thymine or uracil the coding element contains adenine.

5

The complementing element may be a single nucleobase. In the generation of a library, this will allow for the incorporation of four different functional entities into the template-directed molecule. However, to obtain a higher diversity a complementing element preferably comprises at least two and more preferred at least three nucleotides. Theoretically, this will provide for 4<sup>2</sup> and 4<sup>3</sup>, respectively, different functional entities uniquely identified by the complementing element. The complementing element will usually not comprise more than 100 nucleotides. It is preferred to have complementing elements with a sequence of 3 to 30 nucleotides.

10

15

The building blocks of the present invention can be used in a method for transferring a functional entity to a recipient reactive group, said method comprising the steps of providing one or more building blocks as described above and contacting the one or more building blocks with a corresponding coding element associated with a recipient reactive group under conditions which allow for a recognition between the one or more complementing elements and the coding elements, said contacting being performed prior to, simultaneously with, or subsequent to a transfer of the functional entity to the recipient reactive group.

20

The coding element may comprise one, two, three or more codons, i.e. sequences that may be specifically recognised by a complementing element. Each of the codons may be separated by a suitable spacer group. Preferably, all or at least a majority of the codons of the template are arranged in sequence and each of the codons are separated from a neighbouring codon by a spacer group. Generally, it is preferred to have more than two codons on the template to allow for the synthesis of more complex encoded molecules. In a preferred aspect of the invention the number of codons of the encoding element is 2 to 100. Still more preferred are coding elements comprising 3 to 10 codons. In another aspect, a codon comprises 1 to 50 nucleotides and the complementing element comprises a sequence of nucleotides complementary to one or more of the encoding sequences.

25

30

35

SUBSTITUTE SHEET (RULE 26)

The recipient reactive group may be associated with the encoding element in any appropriate way. Thus, the reactive group may be associated covalently or non-covalently to the coding element. In one embodiment the recipient reactive group is linked covalently to the encoding element through a suitable linker which may be separately cleavable to release the reaction product. In another embodiment, the reactive group is coupled to a complementing element, which is capable of recognising a sequence of nucleotides on the encoding element, whereby the recipient reactive group becomes attached to the encoding element by hybridisation. Also, the recipient reactive group may be part of a chemical scaffold, i.e. a chemical entity having one or more reactive groups available for receiving a functional entity from a building block.

The recipient reactive group may be any group able to cleave the bond between the carrier and the functional entity to release the functional entity. Usually, the reactive group is nucleophilic, such as a hydroxyl, a thiol, an amine etc. A preferred recipient reactive group is an amine group. The nucleophile usually attacks the atom of the functional entity connected to the oxygen attached to the nitrogen ring member of the carrier. When the functional entity is attached to said oxygen through a group  $X=V$ , the nucleophile attacks the X atom, thereby causing the carrier group to be a leaving group of the reaction, transferring the  $X(=V)$ -Functional entity precursor to the recipient. The chemical structure formed has, in the event the nucleophilic group is an amine attached to a scaffold, the general formula:



In which


X = -C-, -S-, -P-, -S(O)-, -P(O)-, and

V = O, S, NH, N-C<sub>1</sub>-C<sub>6</sub> alkyl, and R is as previously defined.


In a preferred aspect X is C and V is O.

The conditions which allow for transfer to occur are dependent upon the receiving reactive group. Below various examples of the conditions for a transfer to occur are depicted together with the reaction products formed.

#### A. Acylating building blocks - principle




#### B. Pyrazolone formation by reaction of hydrazines with $\beta$ -Ketoesters




14

### C. Isoxazolone formation by reaction of hydroxylamines with $\beta$ -Ketoesters




### D. Pyrimidine formation by reaction of thioureas with $\beta$ -Ketoesters




5

### E. Pyrimidine formation by reaction of ureas with Malonates



10


### F. Coumarine or quinolinon formation by a Heck reaction followed by a nucleophilic substitution




15

15

### G. Diketopiperazine formation by reaction of Amino Acid Esters



### J. Hydantoin formation by reaction of Urea and $\alpha$ -substituted Esters




5

The present building blocks may be prepared in accordance with a variety of chemical synthesis schemes. Generally, a complementing element containing a thiol group is provided. In the event, the complementing element is a oligonucleotide, the thiol may be provided during the synthesis of the oligonucleotide by incorporating a suitable nucleotide derivative. When a oligonucleotide comprising a thiol group is desired, a variety of commercial nucleotide derivatives are available, e.g. the C6 S-S thiol modifier (obtainable from Glen Research cat. # 10-1936-90), which may be incorporated using the standard protocol of the phosphoramidite synthesis.

10

According to a first synthesis scheme the building block can be prepared using the step



15

The thiol oligonucleotide is reacted with the N-hydroxymaleimide-functional entity derivative via a Michael addition, whereby the SH group is added to the double bond of the maleimide.


20



According to a second synthesis scheme, the building blocks can be prepared in two step:

#### Error! Reference source not found.

The thiol oligonucleotide is reacted with N-hydroxymaleimide via a Michael addition, whereby the SH group is added to the double bond of the maleimide forming an intermediate oligonucleotide derivative which is reacted further with a functional entity connected to a leaving group (Lg). Preferred leaving groups are



According to a preferred aspect of the invention the building blocks are used for the formation of a library of compounds. The complementing element of the building block is used to identify the functional entity. Due to the enhanced proximity between reactive groups when the complementing entity and the encoding element are contacted, the functional entity together with the identity programmed in the complementing element is transferred to the encoding element associated with recipient reactive group. Thus, it is preferred that the sequence of the complementing element is unique in the sense that the same sequence is not used for another functional entity. The unique identification of the functional entity enable the possibility of decoding the encoding element in order to determine the synthetic history of the molecule formed. In the event two or more functional entities have been transferred to a scaffold, not only the identity of the transferred functional entities can be determined. Also the sequence of reaction and the type of reaction involved can be determined by decoding the encoding element. Thus, according to a preferred embodiment of the invention, each different member of a library comprises a complementing element having a unique sequence of nucleotides, which identifies the functional entity.

#### Brief description of the drawings

Fig. 1 shows to setups for functional entity transfer.

Fig. 2 shows examples of specific base pairing.

Fig. 3 shows examples of non-specific base-pairing

Fig. 4 shows examples of backbones.

Fig. 5 discloses the results of example 7.

Fig. 6 discloses the results of example 8.

#### Detailed Description of the Invention

A building block of the present invention is characterized by its ability to transfer its functional entity to a receiving chemical entity. This is done by forming a new covalent bond between the receiving chemical entity and cleaving the bond between the carrier moiety and the functional entity of the building block.

Two setups for generalized functional entity transfer from a building block are depicted in figure 1. In the first example, one complementing element of a building block recognizes a template carrying another functional entity, hence bringing the functional entities in close proximity. This results in a reaction between functional entity 1 and 2 forming a covalent bond between these concurrent with the cleavage of the bond between functional entity 2 and its linker. In the second example, a template brings together two building blocks resulting in functional entity transfer from one building block to the other.

In a library synthesis, several building blocks are mixed in a reaction vessel and the added templates ensure that the building blocks - consequently the functional entities - are combined in the desired manner. As several building blocks are employed at the same time, the use of *in situ* generated building blocks is disfavoured for practical reasons.

Building blocks for library synthesis should possess the necessary reactivity to enable the transfer of the functional entity but should also be stable enough to endure storage and the conditions applied during library synthesis. Hence fine tuning of the reactivity for a particular building block is vital. The reactivity of a building block depends partly on the characteristics of the functional entity and the characteristics of the carrier. E.g. a highly reactive functional entity attached to a highly reactive carrier would form a building block that may be susceptible to hydrolysis during the library synthesis thus preventing successful transfer of one functional entity to another. Further, if transfer of a functional entity precursor is faster than coding element - complementing element recognition unspecific reactions may result.



R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring, or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

5

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, OC(=O)R<sup>18</sup>, OC(=O)OR<sup>18</sup>,

10 OC(=O)NR<sup>18</sup>R<sup>19</sup>, SR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>OR<sup>19</sup>, NR<sup>18</sup>NR<sup>19</sup>R<sup>20</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, P(=O)(OR<sup>18</sup>)OR<sup>19</sup>, C(=O)R<sup>18</sup>, C(=NR<sup>18</sup>)R<sup>19</sup>, C(=NOR<sup>18</sup>)R<sup>19</sup>, C(=O)OR<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup>, C(=O)NR<sup>18</sup>NR<sup>19</sup>R<sup>20</sup>, C(=NR<sup>18</sup>)NR<sup>19</sup>R<sup>20</sup>, C(=NOR<sup>18</sup>)NR<sup>19</sup>R<sup>20</sup> or R<sup>21</sup>,

15

wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring, or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

20

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl, optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=NOR<sup>18</sup>)R<sup>19</sup>, C(=O)OR<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup> or R<sup>21</sup>,

25

wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring, or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

30

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, aziridinyl, azetidyl, pyrrolidinyl, piperidinyl, morpholinyl,

35

phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup>, C(=O)OR<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup> or R<sup>21</sup>,

wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring, or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

10

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=NOR<sup>18</sup>)R<sup>19</sup>, C(=O)OR<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup> or R<sup>21</sup>,

15

wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring, or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

20

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, aziridinyl, azetidyl, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=NOR<sup>18</sup>)R<sup>19</sup>, C(=O)OR<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup> or R<sup>21</sup>,

25

wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring, or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

35

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, phenyl, naphthyl, thieryl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)OR<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup> or R<sup>21</sup>, wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup> or R<sup>21</sup>, wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, thieryl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup> or R<sup>21</sup>, wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl or heteroaryl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)OR<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup> or R<sup>21</sup>, wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thieryl, furyl, pyridyl, quinolinyl or isoquinolinyl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, aziridinyl, azelidinyl, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)OR<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup> or R<sup>21</sup>, wherein,

R<sup>18</sup>, R<sup>19</sup>, R<sup>20</sup> and R<sup>21</sup> independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thieryl, furyl, pyridyl, quinolinyl or isoquinolinyl and wherein R<sup>18</sup> and R<sup>19</sup> may together form a 3-8 membered heterocyclic ring or R<sup>18</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring or R<sup>19</sup> and R<sup>20</sup> may together form a 3-8 membered heterocyclic ring,

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is H, phenyl, naphthyl, thieryl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN, CF<sub>3</sub>, OR<sup>18</sup>, S(=O)<sub>2</sub>R<sup>18</sup>, S(=O)<sub>2</sub>NR<sup>18</sup>R<sup>19</sup>, NO<sub>2</sub>, NR<sup>18</sup>R<sup>19</sup>, NR<sup>18</sup>C(=O)R<sup>19</sup>, NR<sup>18</sup>C(=O)OR<sup>19</sup>, NR<sup>18</sup>C(=O)NR<sup>19</sup>R<sup>20</sup>, C(=O)R<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)OR<sup>18</sup>, C(=O)NR<sup>18</sup>R<sup>19</sup>, C(=O)NR<sup>18</sup>OR<sup>19</sup> or R<sup>21</sup>, wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein  $R^{18}$  and  $R^{19}$  may together form a 3-8 membered heterocyclic ring or  $R^{18}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring.

5

In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,

10

$S(=O)_2R^{18}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{18}R^{19}$ ,  $NO_2$ ,  $NR^{18}R^{19}$ ,  $NR^{18}C(=O)R^{19}$ ,  $NR^{18}C(=O)OR^{19}$ ,  $NR^{18}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{18}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{18}R^{19}$ ,  $C(=O)NR^{18}OR^{19}$  or  $R^{21}$ ,

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein  $R^{18}$  and  $R^{19}$  may together form a 3-8 membered heterocyclic ring or  $R^{18}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring.

20

In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,  $S(=O)_2R^{18}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{18}R^{19}$ ,  $NO_2$ ,  $NR^{18}R^{19}$ ,  $NR^{18}C(=O)R^{19}$ ,  $NR^{18}C(=O)OR^{19}$ ,  $NR^{18}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{18}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{18}R^{19}$ ,  $C(=O)NR^{18}OR^{19}$  or  $R^{21}$ ,

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl and wherein  $R^{18}$  and  $R^{19}$  may together form a 3-8 membered heterocyclic ring or  $R^{18}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring.

30

In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is H, methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents se-

35

lected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,  $S(=O)_2R^{18}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{18}R^{19}$ ,  $NO_2$ ,  $NR^{18}R^{19}$ ,  $NR^{18}C(=O)R^{19}$ ,  $NR^{18}C(=O)OR^{19}$ ,  $NR^{18}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{18}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{18}R^{19}$ ,  $C(=O)NR^{18}OR^{19}$  or  $R^{21}$ ,

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, methyl, ethyl, propyl or butyl and wherein  $R^{18}$  and  $R^{19}$  may together form a 3-8 membered heterocyclic ring or  $R^{18}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring.

5

In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is H, aziridinyl, azetidyl, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,  $S(=O)_2R^{18}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{18}R^{19}$ ,  $NO_2$ ,  $NR^{18}R^{19}$ ,  $NR^{18}C(=O)R^{19}$ ,  $NR^{18}C(=O)OR^{19}$ ,  $NR^{18}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{18}$ ,  $C(=O)NR^{18}R^{19}$ ,  $C(=O)NR^{18}OR^{19}$  or  $R^{21}$ ,

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, methyl, ethyl, propyl or butyl and wherein  $R^{18}$  and  $R^{19}$  may together form a 3-8 membered heterocyclic ring or  $R^{18}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring.

20

In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is H, phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,  $S(=O)_2R^{18}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{18}R^{19}$ ,  $NO_2$ ,  $NR^{18}R^{19}$ ,  $NR^{18}C(=O)R^{19}$ ,  $NR^{18}C(=O)OR^{19}$ ,  $NR^{18}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{18}$ ,  $C(=O)NR^{18}R^{19}$ ,  $C(=O)NR^{18}OR^{19}$  or  $R^{21}$ ,

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, methyl, ethyl, propyl or butyl and wherein  $R^{18}$  and  $R^{19}$  may together form a 3-8 membered heterocyclic ring or  $R^{18}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring.

30

In still another embodiment,

- $R^{17}$  and  $R^{24}$  independently is H, phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{19}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ , wherein,
- $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, methyl, ethyl, propyl or butyl and wherein  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring,

10

In still another embodiment,

- $R^{17}$  and  $R^{24}$  independently is H, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{19}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ , wherein,

15

- $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, methyl, ethyl, propyl or butyl and wherein  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring or  $R^{19}$  and  $R^{20}$  may together form a 3-8 membered heterocyclic ring,

20

In still another embodiment,

- $R^{17}$  and  $R^{24}$  independently is methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{19}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ , wherein,

30

$R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

In still another embodiment,

- $R^{17}$  and  $R^{24}$  independently is aziridinyl, azetidiny, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{19}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ , wherein,

5

$R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

In still another embodiment,

- $R^{17}$  and  $R^{24}$  independently is phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{19}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ , wherein,

15

$R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

In still another embodiment,

- $R^{17}$  and  $R^{24}$  independently is phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{19}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ , wherein,

25

$R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

In still another embodiment,

- $R^{17}$  and  $R^{24}$  independently is thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{19}$ ,  $S(=O)_2R^{19}$ ,  $S(=O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{19}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ , wherein,

35

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl,

5 In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is methyl, ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,  $S(O)_2R^{18}$ ,  $S(O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{18}$ ,  $NR^{19}C(=O)OR^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,

10  $C(=O)R^{18}$ ,  $C(=NOR^{19})R^{18}$ ,  $C(=O)OR^{18}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ ,

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

15 In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is aziridinyl, azetidyl, pyrrolidinyl, piperidinyl or morpholinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,  $S(O)_2R^{18}$ ,  $S(O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{18}$ ,  $NR^{19}C(=O)OR^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{18}$ ,

20  $C(=NOR^{19})R^{18}$ ,  $C(=O)OR^{18}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ ,

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

25 In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is phenyl, naphthyl, thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,  $S(O)_2R^{18}$ ,  $S(O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{18}$ ,  $NR^{19}C(=O)OR^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{18}$ ,  $C(=NOR^{19})R^{18}$ ,  $C(=O)OR^{18}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ ,

30 wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

35 In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is phenyl or naphthyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,  $S(O)_2R^{18}$ ,  $S(O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{18}$ ,

$NR^{19}C(=O)OR^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{18}$ ,  $C(=NOR^{19})R^{18}$ ,  $C(=O)OR^{18}$ ,

5  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ ,

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

10 In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl optionally substituted with one or more substituents selected from the group consisting of F, Cl, CN,  $CF_3$ ,  $OR^{18}$ ,  $S(O)_2R^{18}$ ,  $S(O)_2NR^{19}R^{20}$ ,  $NO_2$ ,  $NR^{19}R^{20}$ ,  $NR^{19}C(=O)R^{18}$ ,  $NR^{19}C(=O)OR^{19}$ ,  $NR^{19}C(=O)NR^{19}R^{20}$ ,  $C(=O)R^{18}$ ,  $C(=NOR^{19})R^{18}$ ,

15  $C(=O)OR^{18}$ ,  $C(=O)NR^{19}R^{20}$ ,  $C(=O)NR^{19}OR^{19}$  or  $R^{21}$ ,

wherein,

$R^{18}$ ,  $R^{19}$ ,  $R^{20}$  and  $R^{21}$  independently is H, phenyl, naphthyl, thienyl, furyl, pyridinyl, quinolinyl or isoquinolinyl,

20 In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is H,  $C_1$ - $C_8$  alkyl,  $C_3$ - $C_7$  cycloalkyl,  $C_3$ - $C_7$  cycloheteroalkyl, aryl or heteroaryl

In still another embodiment,

25  $R^{17}$  and  $R^{24}$  independently is H,

In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is  $C_1$ - $C_8$  alkyl,  $C_3$ - $C_7$  cycloalkyl or  $C_3$ - $C_7$  cycloheteroalkyl,

30 In still another embodiment,

$R^{17}$  and  $R^{24}$  independently is methyl, ethyl, propyl or butyl

In still another preferred embodiment

$R^{17}$  and  $R^{24}$  independently is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl

35

In still another preferred embodiment

R<sup>17</sup> and R<sup>24</sup> independently is aziridinyl, pyrrolidinyl, piperidinyl or morpholinyl

In still another embodiment,

5 R<sup>17</sup> and R<sup>24</sup> independently is aryl or heteroaryl

In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is phenyl or naphthyl

10 In still another embodiment,

R<sup>17</sup> and R<sup>24</sup> independently is thienyl, furyl, pyridyl, quinolinyl or isoquinolinyl

## Experiments


15

All oligos used were prepared by standard phosphoramidite chemistry and purchased from DNA technology, Denmark. The type II compounds used were commercially available from Fluka (4-pentynoic acid cat. no: 77055, 5-hexynoic acid cat. no: 53108 and N-tert-butoxycarbonyl beta-alanine cat. no: 15382). The hexapeptide used as scaffold was synthesised using standard Fmoc chemistry and protected at the N-terminal by acetylation and at the C-terminal by formamide formation. The protected hexapeptide was commercially available from Schaefer-N, Denmark.

20

## Example 1: Preparation of type I compound (method A)

25




N-hydroxymaleimide (4 mmol) was mixed with Et<sub>3</sub>N (4 mmol) in DCM (15 mL) at 0 °C. Acetyl chloride (4 mmol) was added and the reaction mixture was left at rt o/n. DCM (15 mL) was added and the reaction mixture was washed with citric acid (3 x 30 mL), NaHCO<sub>3</sub> (2 x 30 mL) and NaCl aq. (30 mL). The organic phase was dried

30

over MgSO<sub>4</sub> and evaporated *in vacuo* to afford acetic acid 2,5-dioxo-2,5-dihydropyrrol-1-yl ester in 41% yield. <sup>1</sup>H NMR (CDCl<sub>3</sub>): 6.74 (s, 2H), 2.32 (s, 2H).




**Example 2: Preparation of building blocks (method A)**

5

A dTSS-oligo (10 nmol) is evaporated to dryness *in vacuo*. The oligo is redissolved in DTT (50  $\mu$ l 100 mM) in 100 mM Sodium-phosphate buffer pH 8.0. Incubate at 37 °C for 1h and purify using a micro-spin column equilibrated with Hepes-OH (100 mM, pH 7.5). The HS-oligo is treated with CTAB (50  $\mu$ L, 1 mM) and the mixture is evaporated to dryness *in vacuo*. The HS-oligo obtained is redissolved in DMF (100  $\mu$ L) and treated with compounds of type I (100  $\mu$ l 100 mM in DMF) for 3h at rt. NaOAc (200  $\mu$ l 1 M, pH = 7.5) is added and the reaction mixture is extracted with EtOAc (2 x 300  $\mu$ L). The loaded oligo is finally purified using a micro-spin column equilibrated with Hepes-OH (100 mM, pH 7.5).

15

**Example 3: Preparation of building blocks (method B)**

20

C<sub>6</sub>S-S-oligonucleotides A to D (10 nmol) is evaporated to dryness *in vacuo*.

A: 5'-GCG ACC TGG AGC ATC CAT CGT S

B: 5'-GAG CAT CCA TCG S

C: 5'-GAC GAG CAT CCA TCG S

D: 5'-CTA GGG ACG AGC ATC CAT CGS

25

S = Thiol C6 SS modifier (Glen# 10-1936)

The oligo is redissolved in DTT (50  $\mu$ l 100 mM) in 100 mM Sodium-phosphate pH 8.0. Incubate at 37 °C for 1h and purify using a micro-spin column equilibrated with Hepes-OH (100 mM, pH 7.5). NHM (50  $\mu$ l 100 mM) in Hepes-OH (100mM, pH 7.5) is added to the obtained HS-oligo and the mixture is incubated at 25°C for 2h. The oligo-S-NHS is then purified using a Microspin columns equilibrated in MS-grade H<sub>2</sub>O and analysed by ES-MS.

5

A: MS (calc): 6723.52; MS (found): 6723.21

10 B: MS (calc): 3938.75; MS (found): 3937.78


C: MS (calc): 4870.36; MS (found): 4869.42

D: MS (calc): 6435.38; MS (found): 6434.57

Four EDC-activated compounds were prepared by mixing 50  $\mu$ L 100mM of each of the compounds (acetic acid, 4-pentynoic acid, N-tert-butoxycarbonyl beta-alanine, and 5-hexynoic acid) in DMF with 50  $\mu$ l 100 mM of EDC in DMF and leave the mixture at rt for 30 min before use. Subsequently, each of the oligo-S-NHS (1 nmol) is redissolved in MES-buffer (10  $\mu$ l 100 mM, pH 6) and treated with 10  $\mu$ l of a DMF solution of the EDC-activated compounds. After 1 h the building blocks are purified using a microspin column equilibrated with 100 mM MES pH6 to obtain oligonucleotide A loaded with acetyl, oligonucleotide B loaded with 4-pentynyl (=FE<sub>1</sub>), oligonucleotide C loaded with N-tert-butoxycarbonyl beta-alanyl (=FE<sub>2</sub>), and oligonucleotide D loaded with 5-hexynyl (FE<sub>3</sub>).

25

ES-MS analysis of the loaded oligonucleotides showed the masses of their corresponding oligo-S-NHS-building blocks shown above, due to the presence of piperidine added during analysis.

**Example 4: Preparation of scaffold building blocks**

10 nmol of the amino-oligo was diluted in 160  $\mu$ L 100 mM Hepes-KOH buffer pH 7.5. *N*-Succinimidyl 3-[2-pyridyldithio]-propionamido, SPDP (40  $\mu$ L 20 mM, Pierce cat # 21857) was added and the mixture was incubated for 2 h at 30°C. The oligo was extracted with ethyl acetate (200  $\mu$ L) and purified using micro spin columns equilibrated with 100 mM Hepes-KOH buffer pH 7.5. The hexapeptide CysPheLys-LysLys (10  $\mu$ L 100 mM) was added and the mixture was incubated overnight at 30°C. The oligo was purified by ammoniumacetate precipitation and analysed by ES-MS.

10 MS (calc): 8386.41; MS (found): 8386.57

Used oligo:

E: 5'-X CGA TGG ATG CTC GTC CCT AGA YZ


15 X = 5'-amino modifier C6 (Glen# 10-1926)

Y = PC spacer (Glen# 10-4913)

Z = Biotin phosphoramidite (Glen# 10-1955)

Example 5: Transfer of a Functional entity

20



Oligonucleotide A loaded with acetyl (250 pmol) was added to oligo F (200 pmol) in 50  $\mu$ L 100 mM MES, pH 6. The mixture was incubated overnight at 25°C. Subsequently, the mixture was purified by gel filtration using a microspin column equilibrated with H<sub>2</sub>O and transfer of the functional entity was verified by electron spray mass spectrometry (ES-MS).

Used oligos:

A: 5'-GCG ACC TGG AGC ATC CAT CGT - acetyl


SUBSTITUTE SHEET (RULE 26)

F: 5'-X ACG ATG GAT CCT CCA GGT CGC

X = 5' Amino-modifier C6 (Glen# 10-1906)

5 MS (calc): 6667.46; MS (found) 6666.64.

Example 6: Transfer of a three different Functional entities



10

Transfer of the first functional entity. Scaffold building block oligo E (400 pmol) was added to oligo B (400 pmol in 25  $\mu$ L MES buffer, pH 6), loaded with 4-pentynyl, and incubated overnight at 15°C. The volume was then adjusted to 50  $\mu$ L and the mixture transferred to a streptavidin-bead slurry (Pharmacia cat #17-5113-01, pre-washed with 100  $\mu$ L MES buffer) and incubated for 10 min at room-temperature, followed by incubation on ice for 10 min. The beads were washed four times with ddH<sub>2</sub>O, resuspended in 100  $\mu$ L 10mM NaOH and incubated for 2 min at room temperature to denature the duplex. The NaOH was removed and the beads were subsequently washed twice with 60°C ddH<sub>2</sub>O. The water was removed and the beads resuspended in 25  $\mu$ L 100 mM MES buffer pH 6.0.

15

20

Transfer of the second functional entity. Oligo C (400 pmol in 25  $\mu$ L MES buffer, pH 6), loaded with *N*-terbutoxycarbonyl beta-alanyl, was added to the beads and the mixture was incubated at 25°C for 2h. The beads were washed four times with ddH<sub>2</sub>O, resuspended in 100  $\mu$ L 10mM NaOH and incubated for 2 min at room temperature to denature the duplex. The NaOH was removed and the beads were subsequently washed twice with 60°C ddH<sub>2</sub>O. The water was removed and the beads resuspended in 25  $\mu$ L 100 mM MES buffer pH 6.0.


25

SUBSTITUTE SHEET (RULE 26)

Transfer of the third functional entity: Oligo D (400 pmol in 25  $\mu$ l MES buffer, pH 6), loaded with 5-hexynyl, was added to the beads and the mixture was incubated at 25°C for 2h. The beads were washed four times with ddH<sub>2</sub>O, resuspended in 100  $\mu$ l 10mM NaOH and incubated for 2 min at room temperature to denature the duplex. The NaOH was removed and the beads were subsequently washed twice with 60°C ddH<sub>2</sub>O. The beads were additionally washed once with 50  $\mu$ l MES buffer and twice with 50  $\mu$ l water. The beads were resuspended in 25  $\mu$ l ddH<sub>2</sub>O and put on UV transilluminator for 2x15 seconds to cleave oligo E from the beads. 25  $\mu$ l 12% ammonia was added and the mixture was incubated for 5 min at 50°C. The sample was spun twice at 5kG, and the supernatant collected. The sample was evaporated to dryness *in vacuo*, and analysed by ES-MS.

MS of the trisubstituted product (calc): 8197.17

MS of the trisubstituted product (found): 8196.80



#### Example 7: Attachment of functional entity to a thio oligo.

The following oligos containing a nucleobase modified with a S-triphenylmethyl protected thio moiety, were synthesised using the conventional phosphoramidite approach:


L: 5'-WCA TTG ACC TGA ACC ATG BTA AGC TGC CTG TCA GTC GGT ACT  
ACG ACT ACG TTC AGG CAA GA

5 M: 5'-WCA TTG ACC TGA ACC ATG TBA AGC TGC CTG TCA GTC GGT ACT  
TCA AGG ATC CAC GTG ACC AG


W was incorporated using the commercially available thiol modifier phosphoramidite (10-1926-90 from Glen research). B is an internal biotin incorporated using the commercially available phosphoramidite (10-1953-95 from Glen research).

To make an SH group available for further reaction, the S-triphenylmethyl protected thio oligo (10 nmol) was evaporated *in vacuo* and resuspended in TEAA buffer (200  $\mu$ l of a 0.1M solution, pH=6.4). AgNO<sub>3</sub> (30  $\mu$ l of a 1 M solution) was added and the mixture was left at room temperature for 1-2 hours. DTT (46  $\mu$ l of a 1M solution) was added and left for 5-10 minutes. The reaction mixture was spun down (20,000 G for 20 minutes) and the supernatant was collected. The solid was extracted with additional TEAA buffer (100  $\mu$ l of a 0.1 M solution, pH=6.4). The pure thio oligo was obtained by conventional EtOH-precipitation.

The L oligo was subsequently reacted with the compound




forming a building block able to transfer an acetyl group to a nucleophilic group like an amine, and the M oligo was reacted with the compound



forming a building block capable of transferring a 3-tert-butoxycarbonylamino-butanyl group to a nucleophilic recipient group.

The reaction may be represented by the reaction scheme:


30



General procedure: The thio oligo (1 nmol) was dried *in vacuo* and treated with the NHS compound shown above in dimethylformamide (50  $\mu$ l of a 0.1 M solution) and left o/n at rt. The thio oligo was spun down (20,000 G for 10 minutes) and the supernatant removed. Dimethylformamide (1 mL) was added and the loaded thio oligo was spun down (20,000 G for 10 minutes). The dimethylformamide was removed and the loaded thio oligo was resuspended in TEAA buffer (25  $\mu$ l of a 0.1M solution, pH=6.4) and analysed by HPLC.

10

The functional entities were transferred to a amino oligonucleotide according to the scheme:



15

General procedure: The template oligo 5'-

BTCTTGCCCTGAACGTAGTCGTAGTCGATCCGCGTTACCGAGAGCTGGATGCTC  
GACAGGTCCCGATGCAATCCAGAGGTCTG (1 nmol) was mixed with the oligos (L or M) loaded with a functional entity (1 nmol) and amino oligo O in hepes-buffer (20  $\mu$ l of a 100 mM HEPES and 1 M NaCl solution, pH=7.5) and water (added to a final volume of 100  $\mu$ l). The oligos were annealed to the template by heating to 50 °C and cooled (-2 °C/30 second) to 30 °C. The mixture was then left o/n at a fluctuating temperature (10 °C for 1 second then 35 °C for 1 second). The oligo complex was attached to streptavidine by addition of streptavidine beads (100  $\mu$ l, prewashed with 2x1 mL 100 mM hepes buffer and 1M NaCl, pH=7.5). The beads were washed with hepes buffer (1mL). The amino oligo was separated from the streptavidine bound complex by addition of water (200  $\mu$ l) followed by heating to 70 °C for 1

minute. The water was transferred and evaporated *in vacuo*, resuspended in TEAA buffer (45  $\mu$ l of a 0.1 M solution) and product formation analysed by HPLC (see Figure 5).

5. Figure 5 shows the transfer of functional entities to an oligo containing a modified nucleobase with an amino group.

A) The top chromatogram show the reference amino oligo O: 5'-GAC CTG TCG AGC ATC CAG CTT CAT GGC TGA GTC CAC AAT GZ. Z contain the modified nucleobase with an aminogroup, incorporated using the commercially available amino modifier C6 dT phosphoramidite (10-1039-90 from Glen research).

B) The middle chromatogram show the streptavidine purified amino oligo O after partial transfer of a acetyl group from oligo L.

C) The bottom chromatogram show the streptavidine purified amino oligo O after the complete transfer of the more lipophilic 3-tertbutoxycarbonylamino-butanyl.

15 The following gradient was used in the obtainment of the chromatograms: 0-3 minutes 100% A then 15% A and 85% B from 3-10 minutes.

The experiment where the template oligo was omitted showed no non-templated product formation. The results indicate that the efficiency of the templated synthesis was 80-100%. The reason for less than 100% efficiency was probably due to hydrolytic cleavage of the functional entity.

#### Example 8: Simultaneous transfer of two functional entities


25 The following oligo containing a nucleobase modified with a carboxylic acid moiety, was synthesised using the conventional phosphoramidite approach:

H: 5'-GAC CTG TCG AGC ATC CAG CTT CAT GGS AAT TCC TCG TCC ACA  
ATG XT

30

X was incorporated using the commercially available carboxy-dT phosphoramidite (10-1035-90 from Glen research).

The modified oligo was provided with a trisamine scaffold according to the scheme:



- 5 Procedure: The modified oligo (1 nmol) was mixed with water (100  $\mu$ L), hepes buffer (40  $\mu$ L of a 200 mM, pH=7.5), NHS (20  $\mu$ L of a 100 mM solution), EDC (20  $\mu$ L of a freshly prepared 1 M solution) and the tetraamine tetrakis(aminomethyl)methane tetrahydrochloride (20  $\mu$ L of a 100 mM solution). The reaction mixture was left *o/n* at room temperature. The volume was reduced to 60  $\mu$ L by evaporation *in vacuo*. The pure oligo was obtained by addition of  $\text{NH}_3$  conc. (20  $\mu$ L) followed by HPLC purification. It was possible to isolate a peak after approximately 6 min using the following gradient: 0-3 minutes 100% A then 15% A and 85% B from 3-10 minutes then 100% B from 10-15 minutes then 100% A from 15-20 minutes. A = 2% acetonitrile in 10 mM TEAA and B = 80% acetonitrile in 10 mM TEAA.

15

The following oligos containing a nucleobase modified with a *S*-triphenylmethyl protected thio moiety, was synthesised using the conventional phosphoramidite approach:

20 K: 5'-WCA ITG ACC TGT CTG CCB TGT CAG TCG GTA CTG TGG TAA CGG  
GGA TCG ACC T

L: 5'-WCA ITG ACC TGA ACC ATG BTA AGC TGC CTG TCA GTC GGT ACT  
ACG ACT ACG TTC AGG CAA GA


25 W was incorporated using the commercially available thiol modifier phosphoramidite (10-1926-90 from Glen research). B is an internal biotin incorporated using the commercially available phosphoramidite (10-1953-95 from Glen research).

30 To make an SH group available for further reaction, the *S*-triphenylmethyl protected thio oligo (10 nmol) was evaporated *in vacuo* and resuspended in TEAA buffer (200

$\mu$ L of a 0.1M solution, pH=6.4).  $\text{AgNO}_3$  (30  $\mu$ L of a 1 M solution) was added and the mixture was left at room temperature for 1-2 hours. DTT (46  $\mu$ L of a 1M solution) was added and left for 5-10 minutes. The reaction mixture was spun down (20,000 G for 20 minutes) and the supernatant was collected. The solid was extracted with additional TEAA buffer (100  $\mu$ L of a 0.1 M solution, pH=6.4). The pure thio oligo was obtained by conventional EtOH-precipitation.

5

The K and L oligo was subsequently reacted with the compound



10

forming a building block capable of transferring the lipophilic *S*-Trityl-4-mercaptobenzoyl group to a recipient nucleophilic group.

15 The transfer reaction is schematically represented below:



20 The template oligo 5'-

BTCCTGCTGAACGTAGTCGTAGTCGATCCCGCTTACCAGAGCTGGATGCTC

GACAGGTCGCGATGCAATCCAGAGGTCG (1 nmol) was mixed with the two thio oligos (K and L) loaded with the same functional entity (S-Trityl-4-mercaptobenzoyl; 1 nmol) and the trisamine oligo H (1 nmol) in hepes-buffer (20  $\mu$ L of a 100 mM hepes and 1 M NaCl solution, pH=7.5) and water (added to a final volume of 100  $\mu$ L). The oligos were annealed to the template by heating to 50 °C and cooled (-2 °C/ 30 second) to 30 °C. The mixture was then left o/n at a fluctuating temperature (10 °C for 1 second then 35 °C for 1 second). The oligo complex was attached to streptavidine by addition of streptavidine beads (100  $\mu$ L, prewashed with 2x1 mL 100 mM hepes buffer and 1M NaCl, pH=7.5). The beads were washed with hepes buffer (1mL). The trisamine scaffold oligo H was separated from the streptavidine bound complex by addition of water (200  $\mu$ L) followed by heating to 70 °C. The water was transferred and evaporated *in vacuo*, resuspended in TEAA buffer (45  $\mu$ L of a 0.1 M solution) and product formation analysed by HPLC (see Figure 6).

15 The HPLC chromatogram shows the transfer of two functional entities to a scaffold oligo with three amino groups.


A) The top chromatogram shows the reference scaffold oligo H.

B) The bottom chromatogram show the streptavidine purified scaffold oligo H after the partial transfer of one (peak at 7.94 minutes) and two (peak at 10.76 minutes) identical functional entities (S-Trityl-4-mercaptobenzoyl). The following gradient was used: 0-3 minutes 100% A, then 15% A, and 85% B from 3-10 minutes then 100% B from 10-15 minutes. A = 2% acetonitrile in 10 mM TEAA and B = 80% acetonitrile in 10 mM TEAA.

25 Due to the lipophilic nature of the functional entities a longer retention time, in the HPLC chromatogram of the scaffolded molecule with two functional entities compared to one functional entity, was observed. The efficiency of the templated synthesis of a scaffolded molecule with the two identical functional entities was about 25% (peak at 10.76 minutes in Figure 6).

### Model Example 1

General route to the formation of acylating building blocks and the use of these:



5 N-hydroxymaleimide (1) may be acylated by the use of an acylchloride e.g. acetylchloride or alternatively acylated in e.g. THF by the use of dicyclohexylcarbodiimide or diisopropylcarbodiimide and acid e.g. acetic acid. The intermediate may be subjected to Michael addition by the use of excess 1,3-propanedithiol, followed by reaction with either 4,4'-dipyridyl disulfide or 2,2'-dipyridyl disulfide. This intermediate (3) may then be loaded onto an oligonucleotide carrying a thiol handle to generate the building block (4). The reaction of this building block with an amine carrying scaffold is conducted as follows:

10 The template oligonucleotide (1 nmol) is mixed with a thio oligonucleotide building block e.g. (4) (1 nmol) and an amino-oligonucleotide scaffold (1 nmol) in hepes-buffer (20  $\mu$ L of a 100 mM hepes and 1 M NaCl solution, pH=7.5) and water (39  $\mu$ L). The oligonucleotides are annealed to the template by heating to 50 °C and cooling

(2 °C/ second) to 30 °C. The mixture is then left o/n at a fluctuating temperature (10 °C for 1 second then 35 °C for 1 second), to yield template bound (5).

5 The above examples are intended to help illustrate the invention, and are not intended to, nor should they be construed to, limit the scope of the invention. Indeed, various modifications of the invention and many further embodiments thereof, in addition to those shown and described herein, will become apparent to those skilled in the art from the full content of this document, including the examples shown above and the references to the scientific a patent literature cited herein. It should further be appreciated that the contents of those cited references are incorporated herein by reference to help illustrate the state of the art. The examples above contain important additional information that can be adapted to the practice of this invention in its various embodiments and the equivalents thereof.


10

# Abbreviations

DCC	N,N'-Dicyclohexylcarbodiimide
DtOH	3,4-dihydro-3-hydroxy-4-oxo-1,2,3-benzotriazine
DIC	Diisopropylcarbodiimide
DIEA	Diethylisopropylamin
DMAP	4-Dimethylaminopyridine
DNA	Deoxyribonucleic Acid
EDC	1-Ethyl-3-(3'-dimethylaminopropyl)carbodiimide-HCl
HATU	2-(1H-7-Azabenzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HBTU	2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HOAt	N-Hydroxy-7-azabenzotriazole
HOBt	N-Hydroxybenzotriazole
LNA	Locked Nucleic Acid
NHS	N-hydroxysuccinimid
OTf	Trifluoromethylsulfonate
OTs	Toluenesulfonate
PNA	Peptide Nucleic Acid
PyBoP	Benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate
PyBroP	Bromo-tris-pyrrolidino-phosphonium hexafluorophosphate
RNA	Ribonucleic acid
TBTU	2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborate
TEA	Triethylamine
RP-HPLC	Reverse Phase High Performance Liquid Chromatography
TBDMS-Cl	Tert-Butyldimethylsilylchloride
5-Iodo-dU	5-Iodo-deoxyriboseuracil
TLC	Thin layer chromatography
(Boc) <sub>2</sub> O	Boc anhydride, di-tert-butyl dicarbonate
TBAF	Tetrabutylammonium fluoride
SPDP	Succinimidyl-propyl-2-dithiopyridyl
CTAB	Cetylammmoniumbromide

## Claims

## 1. A building block of the general formula




5



capable of transferring a functional entity (FE) to a recipient reactive group, wherein the lower horizontal line is a **Complementing Element** identifying the functional entity and the vertical line between the complementing element and the S atom is a **Spacer**.

10

2. The building block of claim 1, wherein the spacer is a valence bond,  $C_1-C_6$  alkylene-A-,  $C_1-C_6$  alkenylene-A-,  $C_2-C_6$  alkynylene-A-, or




said spacer optionally being connected through A to a moiety selected from




20

3. The compound according to claim 1, wherein the **Spacer** is  $C_1-C_6$  alkylene-A-,  $C_1-C_6$  alkenylene-A-,  $C_2-C_6$  alkynylene-A-, or




said spacer optionally being connected through A to a moiety selected from




connecting group;  $R^1$  is selected independently from H,  $C_1-C_6$  alkyl,  $C_1-C_6$  alkenylene-aryl, or aryl; and n and m independently are integers ranging from 1 to 6.

4. The compound according to claim 1, wherein **Spacer** is -A-, a group  $C_1-C_6$  alkylene-A-,  $C_2-C_6$  alkenylene-A-, or  $C_2-C_6$  alkynylene-A- optionally substituted with 1 to 3 hydroxy groups, or




said spacer being connected through A to a linker selected from



20

5. A compound according to claim 4, wherein the **spacer** is  $C_2-C_6$  alkenylene-A-, said spacer being connected through A to a moiety selected from







where A is a valence bond,  $-C(O)NR^2$ ,  $-NR^2-C(O)-$ ,  $-S-$ ,  $-C(O)-O-$  or  $-OP(=O)(O)-$ ; B is a valence bond,  $-S-$ ,  $-NR^2$ , or  $-C(O)-$  and connects to S-C-connecting group; n and m independently are integers ranging from 1 to 10 and

5  $R^2$  is selected independently from H,  $\begin{array}{c} G \\ | \\ \text{---}O\text{---}G\text{---} \\ | \\ \text{---}N\text{---}G \\ | \\ \text{---}G \end{array}$  or  $\begin{array}{c} G \\ | \\ \text{---}N\text{---}G \\ | \\ \text{---}G \end{array}$ , wherein G is H or C<sub>1</sub>-C<sub>6</sub> alkyl; and the spacer is connected to the complementing element through a nucleobase.

6. A compound according to claim 4, wherein the spacer is -A-



10 said spacer being connected through A to a moiety selected from



-B-,  $-(CH_2)_n-B-$ , or where A is a valence bond,  $-NR^2-C(O)-$ ,  $-O-$ , or  $-S-$ ; B is a valence bond,  $-S-$ ,  $-NR^2$ , or  $-C(O)-$  and connects to S-C-connecting group;

n and m independently are integers ranging from 1 to 10 and

15  $R^2$  is selected independently from H,  $\begin{array}{c} G \\ | \\ \text{---}O\text{---}G\text{---} \\ | \\ \text{---}N\text{---}G \\ | \\ \text{---}G \end{array}$  or  $\begin{array}{c} G \\ | \\ \text{---}N\text{---}G \\ | \\ \text{---}G \end{array}$ , wherein G is H or C<sub>1</sub>-C<sub>6</sub> alkyl; and the spacer is connected to the complementing element via a phosphorus group.

20 7. A compound according to claim 6, wherein the phosphorus group is a phosphate or thiophosphate group attached to a 3' or 5' end of a complementing element.

8. The building block according to any of the claims 1 to 7, wherein FE is  $\begin{array}{c} Y \\ | \\ X \end{array}$  where

X = -C-, -S-, -P-, -S(O)-, or -P(O)-;  
Y = O, S, NH, or N-C<sub>1</sub>-C<sub>6</sub> alkyl; and

25 R is H or selected among the group consisting of a C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl, and heteroaryl, said group being substituted with 0-3 R<sup>3</sup>, 0-3 R<sup>5</sup> and 0-3 R<sup>6</sup> or C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>C(O)R<sup>8</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>C(O)OR<sup>8</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>C(O)OR<sup>8</sup> substituted with 0-3 R<sup>3</sup>.

kylene-O-NR<sup>4</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>C(O)R<sup>8</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>C(O)OR<sup>8</sup> substituted with 0-3 R<sup>3</sup>.

where R<sup>4</sup> is H or selected independently among the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl, heteroaryl, said group being substituted with 0-3 R<sup>3</sup> and

5 R<sup>5</sup> is selected independently from -N<sub>3</sub>, -CNO, -C(NOH)NH<sub>2</sub>, -NHOH, -NHNHR<sup>6</sup>, -C(O)R<sup>8</sup>, -SnR<sup>9</sup>, -B(OR<sup>9</sup>)<sub>2</sub>, -P(O)(OR<sup>9</sup>)<sub>2</sub> or the group consisting of C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> alkadienyl said group being substituted with 0-2 R<sup>7</sup>.

where R<sup>6</sup> is selected independently from H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>7</sub> cycloalkyl, aryl or C<sub>1</sub>-C<sub>6</sub> alkylene-aryl substituted with 0-5 halogen atoms selected from -F, -Cl, -Br, and -I; and R<sup>7</sup> is independently selected from -NO<sub>2</sub>, -COOR<sup>8</sup>, -COR<sup>8</sup>, -CN, -OSiR<sup>8</sup>, -OR<sup>8</sup> and -NR<sup>8</sup>.

10 R<sup>8</sup> is H, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, aryl or C<sub>1</sub>-C<sub>6</sub> alkylene-aryl substituted with 0-3 substituents independently selected from -F, -Cl, -

15 NO<sub>2</sub>, -R<sup>3</sup>, -OR<sup>3</sup>, -SiR<sup>3</sup>

R<sup>9</sup> is -O-, -F-, -Cl-, -Br-, -I-, -CN, -NO<sub>2</sub>, -OR<sup>8</sup>, -NR<sup>8</sup>, -NR<sup>8</sup>C(O)R<sup>8</sup>, -NR<sup>8</sup>C(O)OR<sup>8</sup>, -SR<sup>8</sup>, -S(O)R<sup>8</sup>, -S(O)<sub>2</sub>R<sup>8</sup>, -COOR<sup>8</sup>, -C(O)NR<sup>8</sup> and -S(O)<sub>2</sub>NR<sup>8</sup>.

9. A compound according to claim 8, wherein R is H or selected among the group consisting of a C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl, and heteroaryl, said group being substituted with 0-3 R<sup>5</sup> and 0-3 R<sup>6</sup>, or selected among the group consisting of C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>C(O)R<sup>8</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>C(O)OR<sup>8</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>C(O)R<sup>8</sup>, and C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>C(O)OR<sup>8</sup> substituted with 0-3 R<sup>3</sup>.

10. A compound according to claims 8 or 9, wherein R is H or selected among the group consisting of C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, C<sub>3</sub>-C<sub>7</sub> cycloheteroalkyl, aryl, and heteroaryl, said group being substituted with 0-3 R<sup>5</sup> and 0-3 R<sup>6</sup>.

11. A compound according to any of the claims 8 to 10, wherein R is selected among the group consisting of C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>C(O)R<sup>8</sup>, C<sub>1</sub>-C<sub>3</sub> alkylene-NR<sup>4</sup>C(O)OR<sup>8</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>, C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>C(O)R<sup>8</sup>, and C<sub>1</sub>-C<sub>2</sub> alkylene-O-NR<sup>4</sup>C(O)OR<sup>8</sup> substituted with 0-3 R<sup>3</sup>.

12. A compound according to any of the claims 1 to 11, wherein X = C and V = O or S.

13. A compound according to claims 1 to 12, wherein X = C and V = O.

14. A compound according to claims 1 to 13, wherein complementing element is a nucleic acid.

15. A compound according to claims 1 to 14, wherein Complementing element is a sequence of nucleotides selected from the group of DNA, RNA, LNA, PNA, or morpholino derivatives.

16. A library of compounds according to any of the claims 1 to 15, wherein each different member of the library comprises a complementing element having a unique sequence of nucleotides, which identifies the functional entity.

17. A method for transferring a functional entity to a recipient reactive group, comprising the steps of

providing one or more building blocks according to claims 1 to 15,

contacting the one or more building blocks with a corresponding encoding element associated with a recipient reactive group under conditions which allow for a recognition between the one or more complementing elements and the coding elements, said contacting being performed prior to, simultaneously with, or subsequent to a transfer of the functional entity to the recipient reactive group.

18. The method according to claim 17, wherein the coding element comprises one or more coding sequences comprised of 1 to 50 nucleotides and the one or more complementing elements comprises a sequence of nucleotides complementary to one or more of the coding sequences.

19. The method of claims 17 or 18, wherein the recipient reactive group is an amine group, which may be part of a chemical scaffold, and the linkage between the functional entity and the scaffold is of the general chemical structure:

35 Scaffold-NH-X(=V)-R


In which

X = -C-, -S-, -P-, -S(O)-, -P(O)-, and


V = O, S, NH, N-C<sub>1</sub>-C<sub>9</sub> alkyl.

20. The method according to claim 19, wherein X is C and V is O.

21. A process for preparing a building block according to claim 1, comprising the step of



22. A process for preparing a building block according to claim 1, comprising the steps of



where Lg is a leaving group.

23. A process according to claim 18, wherein the leaving group is selected from




Fig. 1




Fig. 2




Fig. 3

I = Inosine




Fig. 4




Fig 5.





Fig. 6



(19) World Intellectual Property  
Organization  
International Bureau



(43) International Publication Date  
25 September 2003 (25.09.2003)

(10) International Publication Number  
WO 2003/078627 A3

(51) International Patent Classification: C07H 21/00

(21) International Application Number:  
PCT/DK2003/000177

(22) International Filing Date: 14 March 2003 (14.03.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
PA 2002 0415 15 March 2002 (15.03.2002) DK  
60364 056 15 March 2002 (15.03.2002) US  
PCT/DK 02/00419 20 June 2002 (20.06.2002) DK  
10175.539 20 June 2002 (20.06.2002) US  
60434.439 19 December 2002 (19.12.2002) US

(71) Applicant (for all designated States except US): NUBEO-  
LUTION A/S (DK/DK); Remnegade 8, 5th floor, DK-2100  
Copenhagen Ø (DK).

(72) Inventors; and

(75) Inventor/Applicants (for US only): GOULIAEV, Alex,  
Hushør (DK/DK); Brendsted 223, DK-3670 Veksø (Sjæl-  
land (DK); PEDERSEN, Henrik (DK/DK); Fredersej  
24, DK-2880 Bagsvaerd (DK); THISTED, Thomas  
(DK/DK); Fjordskrenten 14, DK-3600 Frederikssund  
(DK); LUNDORF, Mikkel, Dybro (DK/DK); Charlote  
Munkvej 31, 2. tv., DK-2400 København NV (DK);  
SAMS, Christian (DK/DK); Jakob Danneferds Vej 4 A,  
1., DK-1973 Fredensborg C (DK); FRANCHI, Thomas

Published:  
— with international search report

(88) Date of publication of the international search report:  
31 December 2003

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: A BUILDING BLOCK CAPABLE OF FUNCTIONAL ENTITY TRANSFER TO NUCLEOPHIL

(57) Abstract: A building block having the dual capabilities of transferring the genetic information e.g. by recognising an encoding  
element and transferring a functional entity to a recipient reactive group is disclosed. The building block can be designed with an  
adjustable transferability taking into account the components of the building block. The building block may be used in the generation  
of a single complex or libraries of different complexes, wherein the complex comprises an encoded molecule linked to an encoding  
element. Libraries of complexes are useful in the quest for pharmaceutically active compounds.

INTERNATIONAL SEARCH REPORT

International Application No  
PCT/DK 03/00177

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C07H21/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 C07H

Documentation searched other than minimum documentation to the extent that such documents are included in the lists searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category\* Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No.

1, 17

A  
WALDER J A ET AL: "COMPLEMENTARY CARRIER  
PEPTIDE SYNTHESIS: GENERAL STRATEGY AND  
IMPLICATIONS FOR PREBIOTIC ORIGIN OF  
PEPTIDE SYNTHESIS"  
PROCEEDINGS OF THE NATIONAL ACADEMY OF  
SCIENCES OF USA, NATIONAL ACADEMY OF  
SCIENCE, WASHINGTON, US,  
vol. 76, no. 1, January 1979 (1979-01),  
pages 51-55, XP000857351  
ISSN: 0027-8424  
the whole document  
-/-

X Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

\* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another claim or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed
- \*T\* later document published after the international filing date in which the subject matter of the application is further developed or in which the invention is further clarified
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents in the art
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

19 September 2003

06/10/2003

Name and mailing address of the ISA  
European Patent Office, P.B. 5818 Paurtman 2  
NL - 2220 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

de Nooy, A

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/DK 03/00177

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category \* Citation of document, with indication where appropriate, of the relevant passages

Relevant to claim No.

A BRUICK R K ET AL: "TEMPLATE-DIRECTED LIGATION OF PEPTIDES TO OLIGONUCLEOTIDES" CHEMISTRY AND BIOLOGY, CURRENT BIOLOGY, LONDON, 68, vol. 3, no. 1, January 1996 (1996-01), pages 49-56, XP000856876 ISSN: 1074-5521.  
the whole document

1, 17

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/DK 03/00177

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:2. ☒ Claims Nos.: 1-23 (in part)  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/2103. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 8.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.☐ No protest accompanied the payment of additional search fees.

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

## Continuation of Box I.2

Claims Nos.: 1-23 (in part)

Present claims 1-23 relate to an extremely large number of possible building blocks. In fact, the claims contain so many options that a lack of clarity within the meaning of Article 6 PCT arises to such an extent as to render a meaningful search of the claims impossible. Consequently, the search has been carried out for those parts of the application which do appear to be clear (and/or concise), namely those parts relating to the building blocks of claim 1 where the functional entity is as defined in claim 8 and where the complementing element is a nucleic acid or a derivative thereof as in claims 14 and 15.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.